

Theoretical Physics
Prof. Ruiz, UNC Asheville
Chapter J Homework. Spinors

HW-J1. Matrix Properties. Given $\vec{A} = A_x \hat{i} + A_y \hat{j} + A_z \hat{k}$, find the matrix $M = \vec{\sigma} \cdot \vec{A}$, where $\vec{\sigma} = \sigma_x \hat{i} + \sigma_y \hat{j} + \sigma_z \hat{k}$, and

$$\sigma_x = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma_y = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \text{and} \quad \sigma_z = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

The calculate each of the following six items:

$$\text{Tr}(M), \quad M^T, \quad M^*, \quad M^\dagger, \quad \det(M), \quad M^{-1}.$$

HW-J2. Pauli Matrices Identity. Given,

$$\vec{A} = A_x \hat{i} + A_y \hat{j} + A_z \hat{k},$$

$$\vec{B} = B_x \hat{i} + B_y \hat{j} + B_z \hat{k},$$

$$\vec{\sigma} = \sigma_x \hat{i} + \sigma_y \hat{j} + \sigma_z \hat{k},$$

show that

$$\vec{\sigma} \cdot \vec{A} \vec{\sigma} \cdot \vec{B} = \vec{A} \cdot \vec{B} I + i \vec{\sigma} \cdot (\vec{A} \times \vec{B}).$$