

Michael J. Ruiz, “A Black Hole in Our Galactic Center,” *The Physics Teacher* **46**, 10 (January 2008), featured on the journal cover and one of the three papers in the issue made available free online at the journal’s website.

The *American Institute of Physics (AIP)* allows authors to post their publications on their personal websites: “You may post the Version of Record (VOR) 12 months after publication, with the credit line and a link to the VOR on AIP Publishing’s site.” For commercial sites such as ResearchGate, authors can upload their preprint or accepted version with a credit line and link. However, they may send privately the VOR to those requesting the work.

<https://publishing.aip.org/resources/researchers/rights-and-permissions/sharing-content-online/>


This publication appeared in *The Physics Teacher* as shown above.  
The link to the VOR is below.

<https://aapt.scitation.org/doi/10.1119/1.2823992>

Courtesy American Institute of Physics

Publications in AIP journals are Copyright © American Institute of Physics.


© A I P



# THE PHYSICS TEACHER

January 2008      Volume 46      Number 1

Published by the American Association of Physics Teachers



# THE PHYSICS TEACHER

January 2008 Volume 46 Number 1

Published by the American Association of Physics Teachers  
Homepage: <http://www.aapt.org/tpt>

## Papers

- 10 A Black Hole in Our Galactic Center

Michael J. Ruiz



- 13 Speed of Sound in Metal Pipes: An Inexpensive Lab  
Elisha Huggins

- 15 Physics First? Survey First!

Boris Korsunsky and Ozymandias Agar

- 20 Escape from the Pipeline:  
Women Using Physics Outside  
Academia

Jill A. Marshall



- 26 Pinhole Glasses

G. Colicchia, M. Hopf,  
H. Wiesner, and D. Zollman



- 29 Physics of Incandescent Lamp Burnout

Paul Gluck and John King

- 36 A Free-Form Power Experiment to Enhance  
Student Creativity

John T. Fons

- 38 Energy and the Confused Student I: Work


John W. Jewett Jr.

- 44 The Biot-Savart Law: From Infinitesimal to  
Infinite

Jeffrey A. Phillips and Jeff Sanny

- 48 Learning Projectile  
Motion with the  
Computer Game  
"Scorched 3D"

John S. Jurcevic



## Columns

- 4 Letters to the Editor

- 6 Guest Editorial

The Evolution Debate, George V. Coyne

- 8 Figuring Physics, Carbon Dioxide Emission,  
Paul Hewitt

- 52 Apparatus for Teaching Physics  
Microcomputer-Based Laboratory for  
Archimedes' Principle and Density of Liquids,  
Concetto Gianino

- 55 Fermi Questions

Larry Weinstein, Editor

- 56 Physics Challenge for Teachers and  
Students, Boris Korsunsky, Editor

- 57 Trick of the Trade

Simple DC Power Supply, George M. Caplan

- 57 From Our Files, Thomas B. Greenslade Jr.,  
Editor

- 58 Little Gems

A Polarizer Demo Using LCDs, Gerd Kortemeyer  
Polarizing Filters Aren't Supposed to Do  
THAT!, Matt Evans

- 60 etcetera..., Albert Bartlett

- 61 For The New Teacher

Tips for Organizing Your Course, Marc Reif

- 63 Websights

Dan MacIsaac, Editor


- 64 Book Reviews

Brain teaser Physics: Challenging Physics  
Puzzlers, by Goran Grimvall, Alan J. DeWeerd,  
reviewer

Inside Back Cover What's Happening!

## This Month's Cover...

Arrows mark the center of the Milky  
Way Galaxy. Observations of stars  
moving in small orbits around  
that point indicate the presence of a  
massive black hole. Mike Ruiz's  
paper, "A Black Hole in Our Galactic  
Center," begins on p. 10 of this  
month's issue.



---

# A Black Hole in Our Galactic Center

**Michael J. Ruiz**, University of North Carolina at Asheville, Asheville, NC

---

An introductory approach to black holes is presented along with astronomical observational data pertaining to the presence of a supermassive black hole at the center of our galaxy. Concepts of conservation of energy and Kepler's third law are employed so students can apply formulas from their physics class to determine the mass of the black hole that resides in the center of the Milky Way.

We first derive the formula that gives the condition for a black hole from conservation of energy. Students encounter the conservation of energy formula where the kinetic energy plus potential energy at one point is set equal to that for a second point: A common application is to find the escape velocity  $v$  from a celestial body of mass  $M$  and radius  $R$  where atmospheric friction is neglected. We send a projectile of mass  $m$  with speed  $v$  from the surface of our large body so that the projectile comes to rest at infinity. The conservation of energy equation then gives:

$$\frac{1}{2}mv^2 - \frac{GMm}{R} = 0 + 0,$$

where we use the gravitational potential energy

$$U(r) = -\frac{GMm}{r}$$

at  $r = R$  (the start of the journey) and  $r = \infty$  (the end of the trip).

Long ago John Michell (1784) and Pierre-Simon Laplace (1796) applied Newtonian concepts to investigate a theoretical celestial body where the escape velocity would be such that light could not escape. Since Newtonian gravitation (1687) and Einstein's general

theory of relativity (1915) are different frameworks, one must be cautious in making analogies between classical situations and black holes of general relativity.<sup>1</sup> But one cannot resist setting  $v = c$  for the escape velocity and arriving at<sup>2,3</sup>

$$R = \frac{2GM}{c^2}.$$

This relation turns out to be the condition for a black hole in general relativity.<sup>4</sup> The parameter  $R$  is called the Schwarzschild radius after Karl Schwarzschild, who discovered the first solution to Einstein's field equations shortly after Einstein proposed the theory of general relativity.<sup>5</sup>

If mass  $M$  is concentrated into a sphere with radius  $R = \frac{2GM}{c^2}$ , the Schwarzschild radius, a black hole forms. Forces become so great that the mass evolves to a point of infinite density called the singularity. However, we expect this central region to be finite when we eventually arrive at a theory that combines general relativity with quantum mechanics.

A spacecraft can orbit the black hole outside the Schwarzschild radius, but should the spacecraft come closer than this special radius, the spacecraft cannot escape. The spacecraft mass then joins the mass of the black hole, and the Schwarzschild radius becomes a trifle larger as a result. Students can calculate the Schwarzschild radius for the Sun and Earth. If one could somehow get all the mass of the Sun into a volume defined by the Schwarzschild radius, you would produce a black hole. This can lead to discussions of

supernovae, natural phenomena whereby black holes form when the most massive of stars die.

Astronomers believe that supermassive black holes lurk in the centers of many galaxies, including our own.<sup>6,7</sup> These extremely large black holes formed ages ago over billions of years as they “ate” matter over long periods of time, growing larger and larger. In recent years, near-infrared studies of stars close to the center of the Milky Way have revealed data from which we can calculate the mass of our central galactic black hole. We will show below how this is accomplished with basic physics, giving introductory students a problem from the frontiers of astronomical research.

R. Schödel et al.<sup>8</sup> have found that the star S2 (S0-2 in Ref. 11) orbits the center of our galaxy with a semi-major axis of about 1000 AU. It comes as close as 120 AU to the galactic center and has an incredibly short orbital period of only 15 years. For a quick comparison, the distance from the Sun to Pluto is 40 AU, and Pluto orbits the Sun once every 250 years. Kepler’s third law enables us to measure the mass of the black hole<sup>9</sup> at the center of our galaxy.

Kepler’s third law<sup>10</sup> for an object of negligible mass  $m$  orbiting a massive body  $M$  is

$$\frac{GM}{4\pi^2} = \frac{a^3}{P^2},$$

where the semi-major axis is  $a$  and the period  $P$ . Dividing our equation by a similar expression for the Sun with mass  $M_0$ , we obtain

$$\frac{M}{M_0} = \frac{(a/a_0)^3}{(P/P_0)^2}.$$

The simplest units are found by setting  $a_0 = 1$  AU and  $P_0 = 1$  year so that  $a$  is then expressed in AU and  $P$  in years. Note that when  $M = M_0$ , we obtain the familiar Kepler form for the solar system in Earth units.<sup>11</sup>

Using  $a = 1000$  AU and  $P = 15$  years, we obtain for the mass of our black hole

$$M = \frac{1000^3}{15^2} M_0 = 4 \times 10^6 M_0,$$

i.e., 4 million solar masses!

Table I contains data for three stars from Ghez et al.,<sup>12</sup> where Schödel’s S2 is included as S0-2. Students can find an estimate of the supermassive black hole for these cases and even employ a spreadsheet. Four more

**Table I. Some Stellar Orbital Parameters from Ghez et al.**

Star	Semi-major Axis $a$ (AU)	Period $P$ (yr)	Closest Approach in AU
S0-2	$919 \pm 23$	$14.53 \pm 0.65$	$122.2 \pm 2.7$
S0-16	$1680 \pm 510$	$36 \pm 17$	$45 \pm 16$
S0-19	$1720 \pm 110$	$37.3 \pm 3.8$	$287 \pm 25$

stars, with larger uncertainties, are given in Ref. 12.

Students should calculate the Schwarzschild radius for a supermassive black hole consisting of 4 million solar masses. How does this compare to 120 AU, the closest approach of S0-2? What about the closest approach made by S0-16? It is also interesting to express these distances in light hours. Students can place an upper observational bound on the spatial volume within which the central mass must reside. A comparison can be made with the concentration of matter in our solar system. Matter confined to a relatively small volume of space enhances the case for a black hole. By considering such questions, students are engaged with the same issues arising from observational data that concern the professional astronomer.<sup>13</sup>

The European Southern Observatory (ESO) has an excellent website<sup>14</sup> with much information about S2. One can view a nice image of the orbit of S2 around the central black hole in the Milky Way as well as a video clip describing its motion. There is also a fairly recent NOVA program, “Monster of the Milky Way,” on the massive black hole in the center of our galaxy. The video of this program can be obtained on DVD.<sup>15</sup>

### Acknowledgment

The author would like to thank his colleague Brian Dennison, UNCA Glaxo Wellcome Distinguished Professor, for stimulating discussions, and the referee for bringing two relevant website links to the author’s attention.

### References

1. A. K. Raychaudhuri, “Black holes and Newtonian physics,” *Gen. Relat. Gravit.* **24**, 281–283 (March 1992) and

- J. Eisenstaedt, "Dark body and black hole horizons," *Gen. Relat. Gravit.* **23**, 75–80 (Jan. 1991).
- R. W. Brehme, "On the radius of black holes," *Am. J. Phys.* **42**, 1039 (Nov. 1974).
  - Many years ago (1979) at a UNC-Chapel Hill workshop, Hendrik van Dam derived this formula for a group of visiting introductory teachers. He facetiously stated that we used the "wrong formulas" (i.e., nonrelativistic) for kinetic energy and potential energy, and our two mistakes canceled to get the correct general-relativistic result!
  - Technically in general relativity, one must interpret the  $r$ -coordinate as that computed from the circumference of a sphere surrounding the center. Due to the curvature of space, the directly measured radial distance between two nested spheres is larger than the difference found using the general-relativistic  $r$ -coordinate. See E. F. Taylor and J. A. Wheeler, *Exploring Black Holes: Introduction to General Relativity* (Addison-Wesley Longman, San Francisco, 2000), Chap. 2, pp. 7-11, 28.
  - For a nice historical treatment of black holes for the general reader, see K.S. Thorne, *Black Holes & Time Warps: Einstein's Outrageous Legacy* (W. W. Norton & Company, New York, 1994).
  - J.M. Pasachoff and A. Filippenko, *The Cosmos: Astronomy in the New Millennium*, 3rd ed. (Thomson Brooks/Cole, Belmont, CA, 2007).
  - M.A. Seeds, *Stars and Galaxies*, 5th ed. (Thomson Brooks/Cole, Belmont, CA, 2007).
  - R. Schödel et al., "A Star in a 15.2-year orbit around the supermassive black hole at the centre of the Milky Way," *Nature* **419**, 694–696 (Oct. 17, 2002).
  - M. Zeilik and S.A. Gregory, *Introductory Astronomy & Astrophysics*, 4th ed. (Saunders College Publishing, Fort Worth, TX, 1998), p. 346. A situation where Kepler's third law needs to be modified is a binary system of black holes orbiting near each other. A nice diagram showing deviations from Kepler's third law in such an extreme case is found in the otherwise advanced paper P. Grandclément, E.ourgoulhon, and S. Bonazzola, "Binary black holes in circular orbits. II. Numerical methods and first results," *Phys. Rev. D* **65**, 044021-12 (2002).
  - For the more general case where the smaller mass is not negligible, see p. 16 of Ref. 9 for a derivation of
 
$$\frac{G(M+m)}{4\pi^2} = \frac{a^3}{P^2}.$$
  - M.J. Ruiz, "Kepler's third law without a calculator," *Phys. Teach.* **42**, 530–532 (Dec. 2004).
  - A.M. Ghez et al., "Stellar orbits around the galactic center black hole," *Astrophys. J.* **620**, 744–757 (2005). The leading author, Andrea Ghez, has published a book and DVD for grade school students to inspire women to become astronomers: A.M. Ghez and J.L. Cohen, *You Can Be a Woman Astronomer* (Cascade Pass, Culver City, CA, 2006).
  - Readers are encouraged to consult the introductory texts of Refs. 6 and 7 for other observations relating to the supermassive black hole in the center of the Milky Way, including historical perspectives.
  - <http://www.eso.org/outreach/press-rel/pr-2002/pr-17-02.html>.
  - <http://www.pbs.org/wgbh/nova/blackhole/>.
- PACS code: 96.00.00


---

**Michael J. Ruiz**, professor of physics at UNCA, specializes in teaching general education courses in light, sound, and astronomy. His innovative e-texts for these courses have been featured on CNN.

**Department of Physics, University of North Carolina at Asheville, Asheville, NC 28804; [ruiz@unca.edu](mailto:ruiz@unca.edu)**

---

Concerned about  
the safety of your students?



Promote safety awareness and encourage safe habits with this essential manual. Appropriate for elementary to advanced undergraduate laboratories.

Members: \$21.95 • Nonmembers: \$27.50  
order online: [www.thephysicsstore.com](http://www.thephysicsstore.com) or call: 301-209-3333